arXiv:1612.07354v1 [math.GN] 21 Dec 2016

$\mathcal{SPM}$ **BULLETIN**

ISSUE 40 (Dec '16)

1. Editor's note

In addition to the beautiful research announcements, this issue includes an announcement of a festive conference dedicated to Selection Principles, celebrating the 60th birthday of Marion Scheepers. This will be an opportunity for all active researchers in the field to exchange new methods and results in the field, and for researchers looking for new, promising research directions to consider this topic in a thorough manner.

With best regards,

Boaz Tsaban, tsaban@math.biu.ac.il
`http://www.cs.biu.ac.il/~tsaban`

2. New conference: Frontiers of Selection Principles

Dear Friends and Colleagues,

We are glad to invite you to the conference Frontiers of Selection Principles, celebrating the 60th birthday of Marion Scheepers. This festive conference will take place at the Dewajtis Campus, Bielanski forest, Warsaw, 21.8–1.9, 2017: 21.8–25.8 tutorial, 26.8 excursion, 27.8–1.9 conference. (This is immediately after the Logic Colloquium and immediately before the Bedlewo meeting on set theoretic topology and analysis.)

Selection principles connect topology, set theory, and functional analysis and make it possible to transport and apply methods from each of these fields to the other ones. It is now one of the most active streams of research within set theory and general topology. This conference will be fully dedicated to selection principles and their applications. It will begin with a one week tutorial, aimed mainly at students and researchers with no prior knowledge of selection principles. This tutorial will provide an overview of the filed, and a detailed introduction to its main methods. Participants with no prior knowledge attending the tutorial will thus fully benefit from the lectures of the second week, and will be able to consider possible directions for research in the field.

The second conference week will consist of invited lectures, by leading experts, on results in the frontiers of selection principles. We recommend that students and participants with expertise in other disciplines attend both weeks of the conference.

The expected accommodation cost is approximately 20EUR per night (full board: 30EUR).

Registration fee: Early registration: 50EUR. Normal registration: 80EUR. Registration fees will help reducing the costs for some students or early career researchers that would otherwise not be able to attend this meeting. Discount possibilities will be provided in future emails.

Tentative list of conference speakers:

Leandro Aurichi	Liljana Babinkostova	Taras Banakh
Angello Bella*	Daniel Bernal Santos	Maddalena Bonanzinga*
Lev Bukovsky	Steven Clontz*	Samuel Da-Silva*
Rodrigo Dias	Rafał Filipow	David Gauld
Ondrej Kalenda	Ljubiša Kočinac*	Adam Krawczyk
Adam Kwela	Andrzej Nowik	SelmaÖzcağ*
Yinhe Peng	Szymon Plewik	Robert Raowski
Masami Sakai	Marion Scheepers	Paul Szeptycki*
Piotr Szewczak	Franklin Tall	Secil Tokgöz
Boaz Tsaban	Tomasz Weiss	Lyubomyr Zdomskyy
Shuguo Zhnag	Ondrej Zindulka	Szymon Zeberski

(* to be confirmed)

Subject to time constraints, students and other participants may have an opportunity to contribute a short lecture on a topic of selection principles.
Special request: Since a part of this workshop is especially accessible to students, we would appreciate your forwarding this message to your students and to other students who may be interested in attending this conference.
The conference is organized by Cardinal Stefan Wyszyski University in Warsaw.
Important: This is the only message distributed widely. Please email Piotr Szewczak (p.szewczak@wp.pl) to be included in the mailing list for details and updates.

On behalf of the organizing committee,

Piotr Szewczak (Cardinal Stefan Wyszynski University and Bar-Ilan University)
Boaz Tsaban (Bar-Ilan University)
Lyubomyr Zdomskyy (Kurt Gdel Research Center)

3. Long announcements

3.1. **On the definability of Menger spaces which are not σ-compact.** Hurewicz proved completely metrizable Menger spaces are σ-compact. We extend this to Čech-complete Menger spaces and consistently to projective Menger metrizable spaces. On the other hand, it is consistent that there is a co-analytic Menger space that is not σ-compact.

`http://arxiv.org/abs/1607.04688`
Franklin D. Tall and Secil Tokgoz

3.2. **Definable versions of Menger's conjecture.** Menger's conjecture that Menger spaces are σ-compact is false; it is true for analytic subspaces of Polish spaces and undecidable for more complex definable subspaces of Polish spaces. For non-metrizable spaces, analytic Menger spaces are σ-compact, but Menger continuous images of co-analytic spaces need not be. The general co-analytic case is still open, but many special cases are undecidable, in particular, Menger topological groups. We also prove that if there is a Michael space, then productively Lindelof Cech-complete spaces are σ-compact. We also give numerous characterizations of proper K-Lusin spaces. Our methods include the Axiom of Co-analytic Determinacy, non-metrizable descriptive set theory, and Arhangel'skii's work on generalized metric spaces.

`http://arxiv.org/abs/1607.04781`
Franklin D. Tall

3.3. **Discrete subsets in topological groups and countable extremally disconnected groups.** It is proved that any countable topological group in which the filter of neighborhoods of the identity element is not rapid contains a discrete set with precisely one onisolated point and

that the existence of a nondiscrete countable extremally disconnected group implies the existence of a rapid ultrafilter.

http://arxiv.org/abs/1608.03546
Evgenii Reznichenko and Ol'ga Sipacheva

3.4. **σ-Ideals and outer measures on the real line.** A *weak selection* on $\mathbb{R}$ is a function $f : [\mathbb{R}]^2 \to \mathbb{R}$ such that $f(\{x, y\}) \in \{x, y\}$ for each $\{x, y\} \in [\mathbb{R}]^2$. In this article, we continue with the study (which was initiated in [ag]) of the outer measures λ_f on the real line $\mathbb{R}$ defined by weak selections f. One of the main results is to show that CH is equivalent to the existence of a weak selection f for which:

$$\lambda_f(A) = \begin{cases} 0 & \text{if } |A| \leq \omega, \\ \infty & \text{otherwise.} \end{cases}$$

Some conditions are given for a σ-ideal of $\mathbb{R}$ in order to be exactly the family $\mathcal{N}_f$ of λ_f-null subsets for some weak selection f. It is shown that there are $2^{\mathfrak{c}}$ pairwise distinct ideals on $\mathbb{R}$ of the form $\mathcal{N}_f$, where f is a weak selection. Also we prove that Martin Axiom implies the existence of a weak selection f such that $\mathcal{N}_f$ is exactly the σ-ideal of meager subsets of $\mathbb{R}$. Finally, we shall study pairs of weak selections which are "almost equal" but they have different families of λ_f-measurable sets.

http://arxiv.org/abs/1608.06210
S. García-Ferreira, A. H. Tomita, and Y. F. Ortiz-Castillo

3.5. **Selectively (a)-spaces from almost disjoint families are necessarily countable under a certain parametrized weak diamond principle.** The second author has recently shown [20] that any selectively (a) almost disjoint family must have cardinality strictly less than $2^{\aleph_0}$, so under the Continuum Hypothesis such a family is necessarily countable. However, it is also shown in the same paper that $2^{\aleph_0} < 2^{\aleph_1}$ alone does not avoid the existence of uncountable selectively (a) almost disjoint families. We show in this paper that a certain effective parametrized weak diamond principle is enough to ensure countability of the almost disjoint family in this context. We also discuss the deductive strength of this specific weak diamond principle (which is consistent with the negation of the Continuum Hypothesis, apart from other features).

Houston Journal of Mathematics 42 (2016), 1031–1046.

http://arxiv.org/abs/1609.05822
Charles J.G. Morgan and Samuel G. Da Silva

3.6. **Notes on the od-Lindelöf property.** A space is od-compact (resp. od-Lindelöf) provided any cover by open dense sets has a finite (resp. countable) subcover. We first show with simple examples that these properties behave quite poorly under finite or countable unions. We then investigate the relations between Lindelöfness, od-Lindelöfness and linear Lindelöfness (and similar relations with 'compact'). We prove in particular that if a T_1 space is od-compact, then the subset of its non-isolated points is compact. If a T_1 space is od-Lindelöf, we only get that the subset of its non-isolated points is linearly Lindelöf. Though, Lindelöfness follows if the space is moreover locally openly Lindelöf (i.e. each point has an open Lindelöf neighborhood).

http://arxiv.org/abs/1206.0722
Mathieu Baillif

3.7. **Relating games of Menger, countable fan tightness, and selective separability.** Steven Clontz, http://arxiv.org/abs/1610.04800

By adapting techniques of Arhangel'skii, Barman, and Dow, we may equate the existence of perfect-information, Markov, and tactical strategies between two interesting selection games. These results shed some light on Gruenhage's question asking whether all strategically selectively separable spaces are Markov selectively separable.

3.8. **Convergence in topological groups and the Cohen reals.** We show that it is consistent to have an uncountable sequential group of intermediate sequential order while no countable such groups exist. This is proved by adding ω_2 Cohen reals to a model of $\diamondsuit$.

http://arxiv.org/abs/1611.04998

Alexander Shibakov

4. Short announcements

4.1. **Hereditarily normal manifolds of dimension > 1 may all be metrizable.** Alan Dow and Franklin D. Tall, http://arxiv.org/abs/1607.04756

4.2. **A long chain of P-points.** Borisa Kuzeljevic and Dilip Raghavan, http://arxiv.org/abs/1607.07188

4.3. **The cofinal structure of precompact and compact sets in general metric spaces.** Aviv Eshed, M. Vincenta Ferrer, Salvador Hernández, Piotr Szewczak, Boaz Tsaban, http://arxiv.org/abs/1607.07669

4.4. **ω^ω-bases in topological and uniform spaces.** Taras Banakh, http://arxiv.org/abs/1607.07978

4.5. **I-convergence classes of sequences and nets in topological spaces.** Amar Kumar Banerjee and Apurba Banerjee, http://arxiv.org/abs/1608.03381

4.6. **The existence of continuous weak selections and orderability-type properties in products and filter spaces.** Koichi Motooka, Dmitri Shakhmatov, Takamitsu Yamauchi, http://arxiv.org/abs/1608.07144

4.7. **Weakly linearly Lindelöf monotonically normal spaces are Lindelöf.** I. Juhász, V. V. Tkachuk, R. G. Wilson, http://arxiv.org/abs/1610.04506

4.8. **On the cardinality of almost discretely Lindelöf spaces.** Santi Spadaro, http://arxiv.org/abs/1611.07267

4.9. **Categorical properties on the hyperspace of nontrivial convergent sequences.** S. Garcia-Ferreira, R. Rojas-Hernandez, Y. F. Ortiz-Castillo, http://arxiv.org/abs/1611.08289

4.10. **First countable and almost discretely Lindelöf T_3 spaces have cardinality at most continuum.** István Juhász, Lajos Soukup, Zoltán Szentmiklóssy, http://arxiv.org/abs/1612.06651

Previous issues. http://front.math.ucdavis.edu/search?&t=%22SPM+Bulletin%22

Contributions and free subscription. Email tsaban@math.biu.ac.il.